

## Precision Measurement of the Cosmic Ray Helium Flux with AMS Experiment

V. CHOUTKO<sup>1</sup>, ON BEHALF OF THE AMS COLLABORATION<sup>2</sup>.

<sup>1</sup> Massachusetts Institute of Technology, MIT, Cambridge, MA 02139, USA.

<sup>2</sup> For the complete list of authors see the AMS Collaboration list in these proceedings.

vitali.choutko@cern.ch


**Abstract:** AMS is a TeV range high energy physics experiment operating since May 2011 on board of the International Space Station on Low Earth Orbit. The precise helium spectrum as measured by AMS is presented in the rigidity range from 2 GV to 3 TV. Below 10 GV the spectrum is modulated by time dependent solar activity. Above 10 GV the spectrum can be parametrized by a single power law spectrum, modulated by solar activity with fitted solar potential value of  $\phi = 0.5 \pm 0.05$  GV. No fine structures were found in the spectrum.

**Keywords:** Cosmic Rays Nuclei, He flux, AMS-02

### 1 Introduction

Numerous measurements of the He flux were done in the past by balloon and satellite experiments [1],[2],[3],[4],[5],[6],[7],[8],[9],[10],[11],[12]. The exact behaviour of the He flux with energy is important for understanding the acceleration mechanism of charged cosmic rays in our Galaxy. In the present paper we report on the measurement of the He flux, which was performed with about a half a billion He events collected by the AMS-02 detector [13] during its first two years of operation on board of the International Space Station.

### 2 AMS Experiment



**Fig. 1:** A Layout of the AMS-02 detector

The layout of the AMS-02 detector is shown in Figure 1. It consists of a permanent magnet with 0.14 T magnetic field central value, 9 planes of silicon tracker, four layers of time of flight scintillator counters (TOF), a transition radiation detector (TRD), an array of anti-coincidence counters inside of the magnet (ACC), a ring imaging Cerenkov detector (RICH) and an electromagnetic calorimeter (ECAL).

The silicon tracker, which can be further subdivided into

an inner tracker, combining layers 2 to 8, and two external layers 1 (L1) and 9 (L9), measures the particle momentum  $p$  per unit charge  $Z$ , or rigidity,  $R = p/Z$  and the sign of its charge by analysing the particle trajectory behaviour in the magnetic field. The trajectory is determined by up to nine 3D coordinate measurements along the tracker layers of  $300 \mu\text{m}$  thick double sided silicon sensors. With a spatial resolution in the bending He particle direction of 6 to  $7 \mu\text{m}$ , the average maximum detectable rigidity (MDR), i.e.  $R$  for which  $\Delta R/R=1$  is estimated to be  $3.2 \pm 0.3$  TV [14].

The charge magnitude  $Z$  can be measured independently by the charged particle energy loss along the particle trajectory by the tracker L1, TRD, inner tracker, TOF, RICH, tracker L9, and ECAL. As an example, the charge resolution of the inner tracker is about 0.08 charge units for He nuclei [15].

The response of the detector was simulated using the GEANT4-4.9.4 package [16]. The effects of energy loss, scattering, electromagnetic and hadronic interactions, the measured detector resolution, together with the precise geometry description were included. In particular, two different models of He nuclear interactions were used, one is Geant4 Ion Light Binary Cascade model, and the other one is DPMJET-II.5 [17] for rigidities above 10 GV.


### 3 Analysis

The incident particle rigidity together with local coordinates and directions was obtained by fitting these parameters to the 3D coordinates measured in L1, at least 4 out of 7 layers of the inner tracker and L9. The velocity of the particle,  $\beta = v/c$ , was determined using the information of the time of flight hits matching the reconstructed track. The velocity resolution for the He particles was  $\Delta\beta/\beta^2 \approx 0.02$  [18].

A downgoing particle was selected as a helium candidate if the determination of its charge magnitude along the trajectory was consistent with that of He. The main potential source of background to the helium sample were protons with wrongly reconstructed charge and/or high  $Z$  ions interacting before tracker Layer 1. Using the independent measurements of the charge magnitude obtained along the particle trajectory, the proton and ion background was estimated to be less than  $10^{-5}$  and  $10^{-3}$  over all rigidity range correspondingly. As an example, Figure 2 shows

the distribution of the measured particle charge in the inner tracker.

To reject events with large scattering, a  $\chi^2 < 10$  cut on the trajectory fitting quality was applied as shown on Fig. 3. This cut eliminated 1 to 2% of He events, while significantly (factor 3 or more) reducing the number of events with wrongly measured rigidity  $R^1$ .



**Fig. 3:** A Track fit quality cut efficiency dependence from rigidity estimated by ECAL energy deposition. The efficiency is constant within 0.5%.

Next, only events with rigidity above the geomagnetic cutoff were selected, namely :

$$R > R_C \times (1.2 + 2\sigma(R_C))$$

where  $R_C$  was the maximum cutoff rigidity for the events in the AMS field of view, calculated in the dipole approximation [19] for each second time interval along the ISS trajectory around the Earth, and  $\sigma(R_C)$  was the measurement uncertainty at that rigidity value.

#### 4 Helium Flux

The differential helium flux was obtained by firstly unfolding[20] the measured rates with the parametrized<sup>2</sup> detector resolution function and then correcting those for the accumulated time and detector acceptance as a function of the particle rigidity and direction. The accumulated time varied with the rigidity and reached 51.2 million seconds for the rigidities above  $\approx 30$  GV, with the average livetime of 0.81 during the two year measurement. The acceptance was determined by the Monte Carlo method using simulated helium events which underwent the same reconstruction and selection procedure as for the data and was found to be nearly rigidity independent for rigidities above 10 GV. The acceptance needed to be further corrected for the trigger efficiency, the latter being studied by Monte Carlo method and *unbiased trigger*<sup>3</sup> events. The trigger efficiency was found to be  $\approx 95\%$  for the rigidities above 20 GeV and up to 99% for the lower rigidities.

The systematic errors on the calculated flux were the following:

- Due to trigger and event reconstruction efficiency variations relating to constantly changing particle

rates and thermal environment. Above  $\approx 20$  GV this systematic error was found to be nearly rigidity independent. Table 1 shows the average contributions.

- Monte Carlo acceptance evaluation. The calculated average contributions are shown in Table 2.
- Unfolding errors, which arised from uncertainty of knowledge of resolution functions obtained by simulation. These errors are bin to bin correlated, and were estimated to be less than 0.5% below 250 GV , while reaching 10% above 2 TV;

Source	Rigidity Range (GV)	Error (%)
Trigger	2-20	0.3
	>20	0.5
Track & velocity fit	2-20	0.5
	>20	0.7
Total	2-20	0.6
	> 20	0.85

**Table 1:** Average systematics of the trigger and reconstruction.


Source	Rigidity Range (GV)	Error (%)
MC Statistics	> 2	0.7
He selection	2-20	0.5
	> 20	0.7
Geomagnetic cutoff	2-20	0.5
	20-30	0.2
	> 30	0
Total of the above	>2	1.0
He interactions	> 2	3.5

**Table 2:** Average systematics of the Monte Carlo corrections.

The estimations of the systematic errors were verified by varying selection criteria, comparing the obtained average fluxes above  $\approx 20$  GV at different magnetic latitudes as well as comparing the details of He interactions between data and simulation events using the charge measurements along the He trajectory as measured by AMS (see Fig.4).

The unfolding errors were estimated by changing the resolution matrix MDR by about 10%, which corresponded to our test beam data extrapolation error to that rigidity, and allowing up to  $1/20 \text{ TV}^{-1}$  shift in the average inverse rigidity measurement, which corresponded to our current knowledge of tracker alignment using electron and positron samples[21, 22].

1. This was verified by comparing the number of events measured with negative rigidity before and after the cut.
2. by the sum of two gaussian functions
3. This trigger required a coincidence of 3 out of 4 TOF counters in different layers with an amplitude over threshold of  $\approx 1/3$  of that of a minimum ionizing particle and is estimated to have an efficiency exceeding 99.9% for Z=2 particles



**Fig. 2:** A Measured by Inner Tracker charge distribution for helium candidate events selected by Tracker L1 and TOF. Arrows show the values of cut applied. The efficiency of the cut is about 99.8%.

## 5 Analysis of the He Flux Spectrum

To investigate the consistency of the spectrum above 10 GV with the power law spectrum, a fit to the measured event counts by a power law spectrum  $\Phi_0/R^\gamma$  modulated with the solar modulation parameter in the force field approximation [23]  $\phi$  was done by folding this spectrum with the detector resolution function, the estimated acceptance and the measurement time in the inverse rigidity range from  $-0.0005$  to  $0.1 \text{ GV}^{-1}$ . The  $\chi^2/n.d.f$  of the fit was 81/76. The fitted solar modulation potential  $\phi$  value amounts to  $\phi = 0.5 \pm 0.05 \text{ GV}$ . Apart from statistical errors, only the rigidity dependent systematic errors ( $\approx 1.3\%$  above 20 GV) were included into the fit. The  $\chi^2/n.d.f$  became 51/60 for the fit being done in the inverse rigidity range from  $-0.0005$  to  $0.03 \text{ GV}^{-1}$ , with no systematic change of the fitting parameters.

## 6 Results


Fig. 5 shows the helium spectrum as measured by AMS, multiplied by the corresponding bin central value<sup>4</sup> in the 2.7 power together with data of previous experiments.

**Acknowledgments:** This work has been supported by persons and institutions acknowledged in [25] as well as by the Italian Space Agency under contracts ASI-INFN I/002/13/0 and ASDC/011/11/1.


## References

- [1] K. Asakimori et al., The Astrophysical Journal, Vol.502, (1998), 278
- [2] R. Bellotti et al., Physical Review D 60, Issue 5, (1999), 1
- [3] W. Menn et al., The Astrophysical Journal, Vol.533, (2000), 281
- [4] J. Alcaraz et al., Phys. Lett. B 494 (2000), 193.
- [5] T. Sanuki et al., The Astrophysical Journal, Vol.545, (2000), 1135.
- [6] M. Boezio et al., Astroparticle Physics, Vol.19, (2003), 583.
- [7] J.P. Wefel, J.H. Adams, H.S. Ahn et al., Proc. 29th International Cosmic Ray Conference, 2005, 3: p. 105-108.
- [8] V.A. Derbina et al., The Astrophysical Journal, Vol.628,(2005), 41
- [9] Y. Shikaze et al. Astroparticle Physics, Vol.28, (2007) 154
- [10] Y.S. Yoon et al., The Astrophysical Journal, Vol.728, (2011), 122
- [11] O. Adriani et al., Science, Vol.332, (2011) 69
- [12] J. Buckley et al., The Astrophysical Journal, Vol.429, (1994),736
- [13] A. Kounine, Int. J. Mod. Phys. E 21, 1230005 (2012).
- [14] P. Zucco et al., proceedings of the ICRC-2013, to be published.

4. which have been chosen following the recommendations in [24]



**Fig. 5:** The AMS Helium spectrum multiplied by the rigidity value in the 2.7 power together with the previous experimental data.



**Fig. 4:** The distribution of the measured charge at lower TOF for the He events with rigidity greater than 20 GV, selected by Tracker L1 tight charge cut. The difference in number of interacted events between data and MC was estimated to be about 3%.

- [20] A. Kondor, Nucl. Inst. Meth. 216 (1983) 177;  
 G. D'Agostini, Nucl. Inst. Meth. A 362 (1995) 487
- [21] S. Haino et al., proceedings of the ICRC-2013, to be published.
- [22] C. Delgado et al., proceedings of the ICRC-2013, to be published.
- [23] R.A Caballero-Lopez, H. J. Moraal. Geophys. Res. 109 (A1), A01101, 2004
- [24] G.D. Lafferty, T.R. Wyatt , Nucl. Instr. Meth. A 355 (1995) 541.
- [25] M. Aguilar et al., PRL110 141102 (2013).

- [15] P. Saouter et al., proceedings of the ICRC-2013, to be published.
- [16] S. Agostinelli et al., NIM A 506 3 (2003) 250-303.
- [17] J. Ranft, Phys. Rev. D 51, 64 (1995).
- [18] Q. Yan et al., proceedings of the ICRC-2013, to be published.
- [19] C. Størmer, The Polar Aurora, Oxford, (1950).